Student Asthma/Allergy Action Plan

(This Page To Be Completed By Parent/Guardian)

Student Name:			Age:		Grade:
School:			Homeroom	Teacher:	
Parent/Guardian:		Phone()	· .	()
_ "_ "					()
Emergency Contact:		Phone()		()
Known Asthma Triggers: Please che	eck the boxes to identify	what can	cause an asthma	episode for	your student.
□ Exercise □ Respiratory/viral □ Pollens □ Animals/dander □ Temperature/weather—humidity, c □ Other—please list:	old air, etc.	Dust/dust Pesticides	:	☐ Grass ☐ Food	
Known Allergy/Intolerance: Please contact with the allergen	check those which apply	and descr	ibe what happe	ns when your	child eats or comes into
Wheat	pinephrine (such as an E r avoid foods, your doct	piPen®) fo	or an allergy, you	ı must provid	le epinephrine at school. If
Medicines: Please list medicines used at Medicine Name	t home and/or to be give Amount/D		ol.	When d	oes it need to be given
I understand that all medici	nos to ha givan et s	chaol m	ust be provide	led by the	narent/guardien
Parent signature:	nes to be given at s		ast be provid		ate:
Reviewed by school nurse/nurse designee:				Da	ate:

Page 2 of 2

Student Asthma/Allergy Action Plan

(This Page To Be Completed By Health Care Provider)

Student Name:	Weight: Date Of Birth:/ / / / (MONTH) (DAY) (YEAR)					
☐ Exercise Pre-Treatment: Administer inhaler (2 inhalations) 15-30 minutes prior to exercise. ☐ PE ☐ Recess						
☐ Albuterol /Levalbuterol HFA inhaler (Proventil, Ventolin, ProAir ☐ Albuterol DPI (ProAir RespiClick)	r. Xopenex)					
Asthma Treatment Give quick relief medication when student has asthma symptoms, such as coughing, wheezing or tight chest. Albuterol /Levalbuterol HFA - 2-4 inhalations (Proventil, Ventolin, ProAir, Xopenex) Use inhaler with valved holding chamber Albuterol DPI (ProAir RespiClick) - 2 inhalations Albuterol inhaled by nebulizer (Proventil, Ventolin, AccuNeb) 63 mg/3 mL 1.25 mg/3 mL 2.5 mg/3 mL Levalbuterol inhaled by nebulizer (Xopenex) 9.31 mg/3 mL 0.63 mg/3 mL 1.25 mg/3 mL May carry & self-administer quick relief medication If symptoms do not improve, quick relief medication can be repeated after 10 minutes Closely Watch the Student after Giving Quick Relief Medication If, after 10 minutes: Symptoms are better, student may return to classroom after notifying parent/guardian If student continues to get worse, CALL 911 & use the Nebraska Schools' Emergency Response to Life-Threatening Asthma or Systemic Allergic Reactions (Anaphylaxis) Protocol	Anaphylaxis Treatment Give epinephrine when student has allergy symptoms, such as hives, with difficulty breathing (chest or neck "sucking in"), lips or fingernails turning blue, or trouble talking (shortness of breath) or vomiting or collapse. □ EpiPen® 0.3 mg □ EpiPen® Jr 0.15 mg □ AUVI-Q® 0.3 mg □ AUVI-Q® Jr. 0.15 mg □ AUVI-q® 0.1 mg □ Other: Lay person flat on back and raise legs. If vomiting or difficulty breathing, let them lie on their side. □ Use epinephrine auto-injector immediately upon exposure to known allergen □ If symptoms do not improve or they return, epinephrine can be repeated after 5 minutes or more □ May carry & self-administer epi auto-injector CALL 911 After Giving Epinephrine & Closely Watch the Student • Notify parent/guardian immediately • Even if student gets better, the student should be watched for more signs & symptoms of anaphylaxis in an emergency facility If student does not get better or continues toget worse, use the Nebraska Schools' Emergency Response to Life-Threatening Asthma or Systemic Allergic Reactions (Anaphylaxis) Protocol					
This Student has the ability to self-manage Student's Health Condition and I authorize Student to self-manage in accordance with this Plan. If medications are self-administered, the school staff must be notified immediately.						
Additional information: (i.e. asthma triggers, allergens)						
Health Care Provider name: (please print)	Phone:					
Health Care Provider signature:	Date:					
Parent signature:	Date:					

Reviewed by school nurse/nurse designee: